## Morphine Sulfate Injection USP\*

## **INSTRUCTIONS FOR USE:**



as part of the risk minimization plan for ®Morphine Sulfate Injection USP.

This material is not intended for promotional use.

To be administered by slow intravenous injection to non-ambulatory patients for the relief of moderate to severe pain by healthcare professional ONLY. ® Morphine Sulfate Injection USP is not indicated as an as-needed analgesic.

- Perform appropriate hand hygiene when preparing any medication.
- Check the order and confirm the right solution and the right patient.
- Remove the aluminum overwrap.
- Firmly squeeze the bag and check for leaks; if leaks found, do not use the bag as sterility may be impaired.
- Perform visual inspection to check for particles; only use clear solution.
- Do NOT dilute.
- Open the puncture port of the bag by twisting the two flaps in opposite directions to break the first flap (see illustration). **NOTE**: The puncture port septum remains closed and sterile right after removal of the tip.
- Follow the infusion set instructions to properly connect to the puncture port.
- Follow the infusion pump instructions to properly connect to infusion set.
  - Program the infusion pump as per the order.
  - Verify the infusion pump programing and make sure you follow your institution core rights for safe medication administration before starting the infusion.

Patients must be monitored for respiratory depression, especially during initiation of @Morphine Sulfate Injection USP or following a dose increase. Please refer to the opioid management procedures implemented by your institution.

Morphine Sulfate Injection USP is for single use only. Discard any unused portion as per your hospital protocol.

\* Please refer to the product monograph for detailed information.

Questions or concerns? / To report adverse events / Contact us

Medinfo: 1844-329-2939 | medinfo@sterinova.com

