


®HYDROmorphone Hydrochloride Injection USP*

INSTRUCTIONS FOR USE:

To be administered by healthcare professional only



This material was developed by Sterinova Inc., as part of the risk minimization plan for ®HYDROmorphone Hydrochloride Injection USP. This material is not intended for promotional use.

- 1 Perform appropriate hand hygiene when preparing any medication.
- 2 Check the order and confirm the right solution and the right patient.
- 3 Remove the aluminum overwrap.
- 4 Firmly squeeze the bag and check for leaks; if leaks found, do not use the bag as sterility may be impaired.
- 5 Perform visual inspection to check for particles; only use clear solution.
- 6 Open the puncture port of the bag by twisting the two flaps in opposite directions to break the first flap (see illustration). Do not dilute.
NOTE: The puncture port septum remains closed and sterile right after removal of the tip.
- 7 Follow the infusion set instructions to properly connect to the puncture port.
- 8 Follow the infusion pump instructions to properly connect the infusion set.
- 9 Program the infusion pump as per the order.
- 10 Verify the infusion pump programming and make sure you follow your institution core rights for safe medication administration before starting the infusion.

Patients should be monitored for respiratory depression, especially during initiation of ®HYDROmorphone Hydrochloride Injection USP or following a dose increase.

® HYDROmorphone Hydrochloride Injection USP is for single use only.

Discard any unused portion as per your hospital protocol.

* Please refer to the product monograph for detailed information.

Questions or concerns? / To report adverse events / Contact us

Medinfo: 1 844-329-2939 | medinfo@sterinova.com

3005 José-Maria-Rosell Ave
Saint-Hyacinthe QC J2S 0J9 Canada
450 252-2520
sterinova.com

 **STERINOVA**